



E-ISSN: 2664-6773

P-ISSN: 2664-6765

Impact Factor: RJIF 5.6

IJCBS 2024; 6(1): 01-04

www.chemicaljournal.org

Received: 03-11-2023

Accepted: 07-12-2023

P Sudha Rani

Lecturer, Department of Microbiology, Y.A. Govt. Degree College (W), Chirala, Bapatla, Andhra Pradesh, India

K Sujatha

Lecturer, Department of Zoology, S.G.A. Govt. Degree College, Yellamanchili, Andhra Pradesh, India

Chinnababu Sanapala

Research Scholar, Department of Marine Living Resources, Andhra University College of Science and Technology, Visakhapatnam, Andhra Pradesh, India

Nagaraju Sampathirao

Research Scholar, Department of Marine Living Resources, Andhra University College of Science and Technology, Visakhapatnam, Andhra Pradesh, India

Charan Sai Suna

P.G. Student, Department of Microbiology, Andhra University College of Science and Technology, Visakhapatnam, Andhra Pradesh, India

Corresponding Author:**P Sudha Rani**

Lecturer, Department of Microbiology, Y.A. Govt. Degree College (W), Chirala, Bapatla, Andhra Pradesh, India

Biological studies of terpenoids and chemical transformations extracted from essential oil of *Cyperus scariosus*

P Sudha Rani, K Sujatha, Chinnababu Sanapala, Nagaraju Sampathirao and Charan Sai Suna

DOI: <https://doi.org/10.33545/26646765.2024.v6.i1a.70>

Abstract

Cyperus scariosus is a potential medicinal herb belonging to the family *Cyperaceae*. The GC-MS analysis of the oil showed cyrene (16.21%) as the major terpene present in it. Cyrene was isolated from the non-polar fraction of the oil using hexane as solvent and characterized using TLC and spectral techniques (IR and ¹H NMR). Cyrene was derivatized to cyrene epoxide by two methods *i.e.* using perbenzoic acid and epichlorohydrin. Further, the oil, its polar fraction (dichloromethane), non-polar fraction (hexane), cyrene and cyrene epoxide were screened for their plant growth regulating property in case of wheat seedlings (HD 2900 and PBW 601). Complete germination was observed above 2.5 µg/mL of all the test fractions in both the cultivars. Moreover, cyrene epoxide was found to be the most effective in enhancing the length of roots and shoots. Seedling vigour index was calculated in order to analyze the enhancement shown by the oil and its various components on the seedlings.

Keywords: *Cyperus scariosus*, wheat plant growth regulation, cyrene, cyrene epoxide

Introduction

Cyperus scariosus is a medicinal herb and exhibits a wide spectrum of biological activities [1], which includes antibacterial and cytotoxic [2], antifungal [3], insecticidal [4] and antinociceptive [5], hepatoprotective [6], hypotensive and bradycardiac [7] properties. *C. scariosus* is widely distributed in India, especially in Chhattisgarh, Bihar, Orissa, West Bengal and Uttar Pradesh [8]. Extensive work has been carried out to show the plant growth regulating properties of terpenoids

A seed is the basic input in agriculture and a high quality seed is the basis of higher agricultural productivity. Quality in seed includes all physiological, biological, pathological and genetic attributes that contribute towards the final yield of a crop. Seed vigour is a concept describing several characteristics, which include rate and uniformity of germination and growth, tolerance to environmental stresses after sowing and retention of performance after storage. Seeds which perform isolated from medicinal plants [9]. Among terpenoids lactones, conjugated sesquiterpenoid lactones, conjugated sesquiterpenoid ketones and their derivatives with α -methylene- γ -lactone moiety have shown enhancement in adventitious root formation in hypocotyl cuttings of *Vigna radiata* [10], bud break in grapevine cuttings [11] and sprouting in sugarcane cuttings [12]. Well in some or all of these aspects are termed as high-vigour seeds [13]. It is generally recognized that the increase in seed dry weight improves seed vigour [14]. Wheat is the world's most widely cultivated food crop. The area sown under this crop is the largest among the area under various other cereal crops. It is eaten in various forms by more than one thousand million human beings and it makes a large contribution to the calories and proteins available to man than from any other food crop [15]. To date no work has been reported on the chemistry and plant growth regulating activity of *C. scariosus* in wheat seeds. The *C. scariosus* oil and the compounds present in it are believed to be responsible for plant growth regulator activity.


Experimental

IR spectra were taken in nujol on Perkin Elmer, Model RX-1 FT-IR spectrophotometer. ¹H NMR spectra were recorded with Bruker AC (400 MHz) as solutions (in CDCl₃) using tetramethylsilane (TMS) as internal reference.

The IR and ^1H NMR spectra were taken at SAIF, Andhra University, Visakhapatnam. The chemical shifts are expressed in δ (ppm) values. GC-MS analysis was obtained from Advanced Instrumentation Research Facility (AIRF), Marine Living Resources, Visakhapatnam, Andhra Pradesh India. The wheat seeds were obtained.

Column chromatography of *Cyperus scariosus* oil into non-polar and polar fractions: *Cyperus scariosus* oil (5 g) was subjected to column chromatography to fractionate it into non-polar and polar fractions. For column chromatography, the column was packed with 300 g of silica gel with 60-120 mesh size activated at 110 °C for 1 h. Oil was dissolved in hexane and then adsorbed on silica gel for 5-10 min. Column was eluted with hexane as solvent to obtain its non-polar fraction which gave one major spot on TLC plate along with other minor spots. Polar (Dichloromethane) fraction gave a number of spots on TLC plate.

Isolation of cyrene from *C. scariosus* oil: Thin layer chromatography of non-polar fraction on silver-nitrate impregnated chromatoplates gave one major spot along with other minor spots. Major spot, after isolation with the help of column chromatography, was identified as cyrene (1) on the basis of spectral analysis when compared with literature ^[16].



Structure of cyrene (1)

Epoxidation of cyrene (1)

Preparation of perbenzoic acid (4): Sodium metal (10.4 g) was added to methanol (200 mL) in small portions in a round bottomed flask fitted with a reflux condenser. The temperature of the flask was maintained at 0 °C using crushed ice. Solution of freshly recrystallized benzoyl peroxide (2) (100 g) was added to chloroform (200 mL, low temp.) in small portions with constant stirring at 0 °C. This solution was added to sodium methoxide in small portions with continuous stirring keeping the temperature 0 °C. The mixture turned milky after 5 min and was poured on crushed ice. The mixture was transferred to a separatory funnel and was allowed to remain undisturbed for 5 min. Aqueous layer was washed with chloroform (2 × 150 mL) to remove the last traces of benzoyl peroxide and methyl benzoate. The aqueous layer containing sodium benzoate (3) was then acidified with 450 mL of ice cold 1 N sulfuric acid (Scheme-I). Finally perbenzoic acid (4) was extracted from the acidified aqueous

layer using chloroform (3 × 150 mL) and dried over anhydrous sodium sulfate. Perbenzoic acid (4) in chloroform solution was stored in refrigerator.

Determination of normality of perbenzoic acid solution

Sodium thiosulfate solution (0.5 N) was taken in a 250 mL conical flask and known amount of chloroform solution of perbenzoic acid (4) was added to it. An equal volume of dilute sulfuric acid and 1 g of KI was added to the flask. Then 1-2 g of Na_2CO_3 was added to the flask and covered with a glass lid for 1-2 min. The mixture was diluted with 10 mL water and few drops of sodium thiosulfate solution were added to it. Starch solution (2 mL) was added to the flask as an indicator to get the blue coloured solution. The solution was finally titrated with sodium thiosulfate solution with vigorous shaking till the blue colour of the solution was discharged ^[17].

1 mL of 0.1 M $\text{Na}_2\text{S}_2\text{O}_3 \equiv 0.0069$ g of perbenzoic acid

Epoxidation using perbenzoic acid

A solution of cyrene (1) (1.5 g) in chloroform (10 mL) was treated with an excess of perbenzoic acid solution in chloroform (Scheme-II). The reaction mixture was stirred continuously while the temperature was maintained at 0 °C with the help of an ice bath. The completion of reaction was checked by thin layer chromatography. The reaction mixture was diluted with chloroform, washed with saturated sodium thiosulfate and sodium bicarbonate solution, finally with brine and then dried over sodium sulfate. Evaporation of solvent afforded a thick yellow liquid (0.8 g), which was chromatographed over silica gel (100 g). Dichloromethane fraction afforded pure cyrene epoxide.

Epoxidation using epichlorohydrin

Cyrene (1) (0.5 g) and epichlorohydrin, $\text{C}_3\text{H}_5\text{ClO}$ (30 mL), were refluxed with continuous stirring for 13 h at 60-65 °C (Scheme-II). On completion of the reaction (TLC), the reaction mixture was filtered and extracted from chloroform. The organic phase was washed with 10% NaOH (50 mL), water and then with brine. The organic phase was then concentrated and column chromatographed to obtain pure epoxy compound. Finally, the oil, its polar and non-polar fraction, cyrene (1) and cyrene epoxide (5) were used for the bio-efficacy studies as plant growth regulator in case of wheat seedlings.

Plant growth regulator studies

Seed germination and seedling establishment studies in

Triticum aestivum: The seeds were germinated in petri-dishes (9 cm) lined with filter paper and moistened with solution of the oil, polar fraction, non-polar fraction, cyrene (1) and cyrene epoxide (5) at different concentrations along with distilled water as control. Ten seeds were planted per petri-dish. The experiment was replicated thrice. The petri-dishes were incubated at 20 ± 2 °C. Petri-dishes were observed daily for 10 days for germination (radical protrusion of 2 mm). Primary root length and shoot length were measured at the end of ten days with a centimeter scale.

At the end of 10 days, seedlings were blotted dry and fresh weight was recorded. For dry weight determination, the seedlings were oven dried at 60 °C for 3 days in an incubator.

Preparation of stock solution

The stock solution (2000 $\mu\text{g}/\text{mL}$) of oil, its polar and non-polar fractions, cyrene (1) and cyrene epoxide (5) was

prepared by dissolving each chemical (20 mg) in 1 mL of Tween 20 (polyoxyethylene sorbitan) and volume was made 10 mL with distilled water. The stock solution of 2000 $\mu\text{g/mL}$ of each compound thus prepared on active ingredient basis and was kept in refrigerator till use. The required dilutions of 25, 20, 10, 5, 2.5, 1 and 0.5 $\mu\text{g/mL}$ were subsequently made from the stock solution by adding distilled water as and when required.

Results and Discussion

C. scariosus oil was obtained from laboratory stock and was reddish brown liquid. The physical properties of the oil were examined such as refractive index (1.39), specific gravity (1.508), viscosity (2.061 poise) and pH (3.3). The essential oil was insoluble in water, sparingly soluble in hexane and completely soluble in acetone and ethanol. Thin layer chromatography of the *C. scariosus* oil showed seven coloured spots having R_f values of 0.95, 0.90, 0.83, 0.70, 0.63, 0.55 and 0.45. Gas chromatography-mass spectrometry (GC-MS) data of isolated *C. scariosus* oil showed the presence of cyrene (1) (18.57%) as the major compound. Other minor compounds present were cyclopropazulen-7-ol, caryophyllene oxide, isolongifolen-5-one, longiverbenone, zierone etc. *C. scariosus* oil was subjected to column chromatography to fractionate it into non-polar (hexane) and polar fractions (dichloromethane). The TLC of non-polar fraction showed one major pinkish-brown spot having $R_f = 0.95$, along with other minor spots. However, the dichloromethane fraction showed six spots having R_f values 0.90, 0.83, 0.70, 0.63, 0.55 and 0.45.

Isolation of cyrene (1) from *Cyperus scariosus* oil

Column chromatography of non-polar (hexane) fraction of *C. scariosus* afforded major spot cyrene (1). The purity of the compound was checked by TLC and the structure was confirmed by spectral analysis.

The ^1H NMR (CDCl_3) spectrum of compound 1 showed singlets at $\delta = 1.62$ ppm (C11-CH_3) which indicated the presence of tetrasubstituted double bond, 0.77 ppm (C13-CH_3) and 0.95 ppm (C14-CH_3). A doublet at 0.80 ppm ($J = 6.4$ Hz) was observed for the $-\text{CH}_3$ group at C12 position. The compound failed to show any signal corresponding to the olefinic protons, thus, it confirmed that the double bond was tetrasubstituted.

The IR analysis of compound 1 showed intense bands at 1382, 1372 cm^{-1} corresponding to C13 and C14 *gem* dimethyl group. A band was observed at 3078 and 1644 cm^{-1} corresponding to C-H stretching and C=C stretching of tetrasubstituted double bond, respectively.

Epoxidation of cyrene

The oxidation of terpenes is an important industrial application as epoxides are used as starting materials in the synthesis of commercially important materials for enhancing fragrance and flavor. Cyrene epoxide (5) was obtained from cyrene (1) by two different methods (Scheme- II). In first method, cyrene (1) was treated with chloroform solution of perbenzoic acid (4) taken in excess with continuous stirring. In another method, cyrene (1) was refluxed with epichlorohydrin for 13 h. In both cases, cyrene epoxide (5) was obtained as a thick yellow liquid. The purity of the compound was checked by thin layer chromatography (Benzene: ethyl acetate, 19:1) having $R_f = 0.65$ and the structure was confirmed by spectroscopic analysis.

The ^1H NMR (CDCl_3) spectrum of compound 5 showed

singlets at $\delta = 1.19$ ppm (C11-CH_3), 0.74 ppm (C13-CH_3) and 0.90 ppm (C14-CH_3). It also showed a doublet at 1.00 ppm ($J = 6.84$ Hz) corresponding to C12-CH_3 group. The chemical shifts observed in case of cyrene epoxide (5) were downfield as compared to those in cyrene (1) because of deshielding occurred due to formation of epoxide ring. Further, the formation of epoxide was determined with the help of IR spectrum. The IR analysis of cyrene epoxide (5) showed the presence of bands at 1277 cm^{-1} corresponding to C-O stretching and 889 and 803 corresponding to C-O bending of epoxy group. The intense bands corresponding to *gem* dimethyl group were found at 1378 and 1316 cm^{-1} .

Evaluation of *Cyperus scariosus* oil and its components for plant growth regulator activity:

After the successful derivatization of cyrene (1) to cyrene epoxide (5), *C. scariosus* oil, its non-polar and polar fractions, cyrene (1) and cyrene epoxide (5) were screened for their plant growth regulation studies. For initial screening of the oil, its fractions, isolated compound and its derivative, we preferred to use those seeds which can give results within 7-10 days e.g. wheat, mung beans etc. Therefore, two different varieties of wheat (*Triticum aestivum*), HD 2967 and PBW 621 were selected. The effect of oil, its non-polar and polar fractions, cyrene (1) and cyrene epoxide (5) were studied on root length, shoot length, fresh weight, dry weight and seedling vigour index.

Observations: The different parameters such as fresh weight, dry weight, root length and shoot length were observed in control and are listed in Table-1.

The following observations were recorded in case of wheat seedlings treated with the oil, its fractions, isolated compound and its derivative:

Germination percentage: Germination percentage of the seeds germinated was calculated using the formula:

Table 1: Physical parameters for HD 2900 and PBW 601 in control

Parameters	HD 2900	PBW 601
Fresh weight	0.223 g	0.211 g
Dry weight	0.059 g	0.052 g
Root Length	9.00 cm	7.00 cm
Shoot Length	12.00 cm	11.00 cm

The germination percentage increased with the increase in concentration for each fraction. At higher concentration (Above 2.5 $\mu\text{g/mL}$) complete germination was observed in all the cases.

Number of roots: The number of roots were found to be four in each case at each concentration of the compounds as well as in control in both cultivars.

Length (shoot and root): Lengths were measured using a centimeter scale. Shoot length was measured from the base till tip of the uppermost leaf. Root length was measured from the base till tip of the longest root. Both shoot and root length increased with the increase in concentration of each compound. Cyrene epoxide (5) was found to be the most active in enhancing shoot and root length at higher concentrations in both varieties. The oil and its polar fraction also showed a good potential in enhancing the shoot and root length.

Fresh weight of seedling: The seedlings were blotted dry on the 10th day of the experiment and fresh weight was recorded. Likewise, fresh weight of the seedlings showed a linear relationship with the concentration of the compounds. The oil, its polar fraction and cyprine epoxide (5) increased the fresh weight of the seedlings more efficiently as compared to cyprine (1) and the non-polar fraction. Cyprine epoxide (5) showed a maximum effect in increasing the weight of the seedlings.

Dry weight of seedling: The seedlings were oven dried at 60 °C for 3 days and then the dry weight was recorded. A similar trend was observed in determination of dry weight of seedling as in case of fresh weight. Cyprine epoxide (5) was found to be the most effective in increasing the dry weight of the seedling.

Seedling vigour index (VI): Seedling vigour index (VI = % germination × total seedling dry weight) also increased with the increase in concentration in all the cases. The results of seedling vigour index showed that the oil, its polar and non-polar fractions, cyprine (1) and cyprine epoxide (5) have improved the performance of the seeds in terms of germination.

Statistical analysis: The statistical analysis for the plant growth regulator activity of the oil, its polar and non-polar fractions, isolated compound cyprine (1) and cyprine epoxide (5) was carried out. The results were analyzed using Factorial CRD. The interactions between concentrations (A), compounds and varieties (C) were analyzed to study the enhancement of root length, shoot length, fresh weight, dry weight and seedling vigour index in case of both the varieties (HD 2967 and PBW 621). The results are as follows:

Wheat cultivar HD 2900: The value of coefficient of variation was found to be 8.2432 for shoot length, 7.7591 for root length and 1.494 for dry weight. The interaction between A and C was found to be significant (least significant difference (LSD) = 0.943) amongst all other factors in case of shoot length. All the interactions *i.e.* between concentrations, compounds and variety with respect to control (*i.e.* without treatment) were significant in root length and dry weight. No significant results were obtained for the interaction between compounds and/or concentrations with respect to the variety for seedling vigour index. Thus, it can be concluded that the effect of concentration was significant with respect to variety as compared to control (*i.e.* without treatment) and all the compounds behave similarly on increasing the shoot length.

Wheat cultivar PBW 601: The coefficient of variation was 8.6021 and the only significant interaction was between A and C (LSD = 0.9375) for shoot length whereas all the interactions *i.e.* between concentrations, compounds and variety with respect to control (*i.e.* without treatment) were significant for root length. The seedling vigour index for the variety was significantly different (LSD = 0.3294) as compared to control (*i.e.* without treatment) and a significant interaction was observed between the concentrations and the variety (LSD = 0.9316).

Conclusion

The epoxidation of cyprine (1) with epichlorohydrin was more convenient as compared to that with perbenzoic acid (4) as well as the yield of the reaction was more in case of

epichlorohydrin. In general, the oil, its polar and non-polar fractions, cyprine and derivatives enhanced the growth of the wheat seedlings. The decreasing order of activity at different concentrations was as follows:

Cyprine epoxide (5) > Polar fraction >
C. scariosus oil > Cyprine (1) > Non-polar fraction

Thus, it can be concluded that there exist a linear relationship between the oil, its polar and non-polar fractions, cyprine and cyprine epoxide (5) with the plant growth regulating activity as with the increase in concentration, the length of the roots and shoots increased. Moreover, it is very interesting to note that cyprine epoxide (5) has been found to be the most active in promoting the root and shoot length and are promising plant growth regulators.

Acknowledgements

The authors are thankful to the Head, Department of Chemical Engineering, Andhra University, Visakhapatnam for providing facilities to carry out this work.

References

1. Utreja D, Sharma P, Ekta Curr. Chem Biol; c2015, 9(2).
2. Rahman MS, Anwar MN Bangladesh J. Microbiol. 2008;25:82.
3. Dixit A, Husain A Fitoterapia. 1984;55:3171.
4. Baskaran J, Dhanasekaran S, Krishnappa K, Anandan A, Elumalai K Int. J. Rec. Sci. Res. 2010;8:183.
5. Alam MA, Jahan R, Rahman S, Das AK, Rahmatullah M Pak. J. Pharm. 2011;24:53.
6. Gilani AH, Janbaz KH Gen. Pharm. 1995;26:627.
7. Gilani AH, Janbaz KH, Zaman M, Lateef A, Tariq SR, Ahmad HR Arch. Pharm. Res. 1994;17:145.
8. Kasana B, Sharma SK, Singh L, Mohapatra S, Singh T. Int. Res. J. Pharm. 2013;4:17.
9. Talwar KK Plant Physiol. Biochem. 1987;14:36.
10. Kalsi PS, Sood VB, Masih AB, Gupta D, Talwar KK. Phytochemistry 1983;22:1387.
11. Kaur S, Cheema SS, Chhabra BR, Talwar KK. Plant Growth Regul. 2002;37:63.
12. Talwar KK, Kanwal RS Sugarcane. 1984;1:13.
13. Black M, Bewley JD. Seed Technology and its Biological Basis. Sheffield Academic Press Ltd, Sheffield, UK; c2000.
14. Hampton JC, Rahman M, Mwakangwale MG, Hill MJ. Seed Sci. Technol. 2005;33:499.
15. Kodesia J. Wheat in India. National Cooperative Printing Press, New Delhi; c1982. p. 1-2.
16. Trivedi B, Motl O, Smolikova J, Sorm F. Tetrahedron Lett. 1964;5:1197.
17. Mann FG, Saunders BC. Practical Organic Chemistry. Orient Longman Press, New Delhi; c2007, 456.